## Biotinylated Human SKP1 Protein

Cat. No. SKP-HM401B



Description	
Source	Recombinant Biotinylated Human SKP1 Protein is expressed from HEK293 His tag at the N-terminus and Avi tag at the C-terminus
	It contains Pro2-Lys163.
Accession	P63208-1
Molecular Weight	The protein has a predicted MW of 21.43 kDa. Due to glycosylation, the protein migrates to 23-28 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.


The S-phase Kinase-Associated Protein 1 (SKP1) is a core component of the SKP1, Cullin 1, F-box protein (SCF) complex, an E3 ubiquitin ligase that serves to poly-ubiquitinate a vast array of protein targets as a signal for their proteasomal degradation, thereby playing a critical role in the regulation of downstream biological

processes.

## **Assay Data**

**Background** 

## Tris-Bis PAGE




Biotinylated Human SKP1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. SKP-HM401B



## **Assay Data**



The purity of Biotinylated Human SKP1 is greater than 95% as determined by SEC-HPLC.