

Human SLAMF1/SLAM/CD150 Protein

Cat. No. SLA-HM1MF

Description

Source	Recombinant Human SLAMF1/SLAM/CD150 Protein is expressed from Expi293 with His tag at the C-terminal. It contains Ala21-Pro237.
Accession	Q13291-1
Molecular Weight	The protein has a predicted MW of 25.4 kDa. Due to glycosylation, the protein migrates to 40-60 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Supplied as 0.22 μm filtered solution in PBS (pH 7.4). Please dilute to the desired concentration according to the concentration of the solution shown on the product label.
Storage	Valid for 12 months from date of receipt when stored at -80°C . Recommend to aliquot the protein into smaller quantities for optimal storage. Please do not repeated freeze-thaw cycles.

Background

The signaling lymphocytic activation molecule (SLAM) family is comprised of nine distinct receptors (SLAMF1 through SLAMF9) that are expressed on hematopoietic cells. The SLAM family receptor function is largely controlled via SLAM associated protein (SAP) family adaptors. The SAP family adaptors consist of SAP, Ewing sarcoma associated transcript (EAT)-2, and EAT-2-related transducer (ERT).


Assay Data

Tris-Bis PAGE



Human SLAMF1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human SLAMF1 is greater than 95% as determined by SEC-HPLC.