## **Human SPARC Protein**

#### Cat. No. SPA-HM101



Description	
Source	Recombinant Human SPARC Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Ala18-Ile303.
Accession	P09486
Molecular Weight	The protein has a predicted MW of 33.8 kDa. Due to glycosylation, the protein migrates to 40-50 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

#### Formulation and Storage


Formulation and Storage		
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.	
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.	
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	

# **Background**

Secreted protein acidic and rich in cysteine (SPARC/osteonectin/BM40) is one of the most abundant noncollagenous protein expressed in mineralized tissues. The capacity of SPARC to influence pathways involved in extracellular matrix assembly such as procollagen processing and collagen fibril formation as well as the capacity to influence osteoblast differentiation and osteoclast activity will be addressed.

## **Assay Data**


#### **Tris-Bis PAGE**



Human SPARC on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

# KAGTUS


## **Assay Data**



The purity of Human SPARC is greater than 95% as determined by SEC-HPLC.

#### **ELISA Data**

Human SPARC, His Tag ELISA 0.05µg Human SPARC, His Tag Per Well



Immobilized Human SPARC, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-SPARC Antibody, hFc Tag with the EC50 of 0.10µg/ml determined by ELISA.