

Cynomolgus SFRP2 Protein

Cat. No. SRP-CM102



Description

Source	Recombinant Cynomolgus SFRP2 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Leu25-Cys295.
Accession	G8F2Z7
Molecular Weight	The protein has a predicted MW of 32.16 kDa. Due to glycosylation, the protein migrates to 33-38 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

As biomarkers, DNA methylation is used to detect colorectal cancer (CRC) and make assessment of CRC prognosis. The published findings showed the association between the methylation of SFRP1, SFRP2, and WIF1, located in the Wnt signaling pathway, and the prognosis of CRC were not consistent. SFRP1, SFRP2, and WIF1 were frequently hypermethylated in CRC tumor tissues. It was apparent that the promoter hypermethylation of SFRP2 and co-hypermethylation of SFRP1 and SFRP2 might be considered as independent prognostic predictors for survival advantage of postoperative CRC patients.


Assay Data

Bis-Tris PAGE



Cynomolgus SFRP2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Cynomolgus SFRP2 is greater than 95% as determined by SEC-HPLC.