

Human SIRP alpha V2/CD172a Protein

Cat. No. SRP-HM1V2

Description

Source	Recombinant Human SIRP alpha V2/CD172a Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Glu31-Arg369.
Accession	AAH26692.1
Molecular Weight	The protein has a predicted MW of 38 kDa. Due to glycosylation, the protein migrates to 55-65 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Signal regulatory protein α (SIRP α) is a regulatory membrane glycoprotein from SIRP family expressed mainly by myeloid cells and also by stem cells or neurons. SIRP α acts as inhibitory receptor and interacts with a broadly expressed transmembrane protein CD47 also called the "don't eat me" signal. Cancer cells highly expressed CD47 that activate SIRP α and inhibit macrophage-mediated destruction.

Assay Data


Tris-Bis PAGE



Human SIRP alpha V2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human SIRP alpha V2 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human SIRP alpha V2, His Tag ELISA

0.05µg Human SIRP alpha V2, His Tag Per Well




Immobilized Human SIRP alpha V2, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-SIRP alpha Antibody, hFc Tag with the EC50 of 4.0ng/ml determined by ELISA (QC Test).

ELISA Data

Human SIRPA V2, His Tag ELISA

0.1µg Human SIRPA V2, His Tag Per Well




Immobilized Human SIRP alpha V2, His Tag at 1µg/ml (100µl/Well) on the plate. Dose response curve for Human CD47, hFc Tag with the EC50 of 0.15µg/ml determined by ELISA.

Blocking Data

Assay Data

Inhibition of Human SIRP alpha V2 and CD47 Binding

0.5µg Human SIRP alpha V2, His Tag Per Well



Serial dilutions of Anti-SIRP alpha Antibody, hFc Tag were added into Human SIRP alpha V2, His Tag : Biotinylated Human CD47 Protein, His Tag binding reactions. The half maximal inhibitory concentration (IC50) is 54.14ng/ml.