Human SIRP alpha V2/CD172a Protein


Cat. No. SRP-HM2V2

ϗͶϲϿ·ႮႽ

| Description | |
|-------------------------|--|
| Source | Recombinant Human SIRP alpha V2/CD172a Protein is expressed from HEK293 with hFc tag at the C-Terminus. |
| | It contains Glu31-Arg369. |
| Accession | AAH26692.1 |
| Molecular Weight | The protein has a predicted MW of 63.9 kDa. Due to glycosylation, the protein migrates to 75-85 kDa based on Tris-Bis PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Tris-Bis PAGE |
| | > 95% as determined by HPLC |
| Formulation and Storage | |
| Formulation | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization. |
| Reconstitution | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water. |
| Storage | -20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |
| Background | |
| | Signal regulatory protein α (SIRP α) is a regulatory membrane glycoprotein from SIRP family expressed mainly by myeloid cells and also by stem cells or neurons.SIRP α acts as inhibitory receptor and interacts with a broadly expressed transmembrane protein CD47 also called the "don't eat me" signal.Cancer cells highly expressed CD47 that activate SIRP α and inhibit macrophage-mediated destruction. |

Assay Data

Tris-Bis PAGE



Human SIRP alpha V2 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%..


SEC-HPLC

Human SIRP alpha V2/CD172a Protein

Cat. No. SRP-HM2V2

Assay Data






The purity of Human SIRP alpha V2 is greater than 95% as determined by SEC-HPLC.




0.1µg Human CD47, His Tag Per Well



Immobilized Human CD47 at 1 μ g/ml (100 μ I/Well). Dose response curve for Human SIRP alpha V2, hFc Tag with the EC50 of 30.6ng/ml determined by ELISA (QC Test).

ELISA Data



Human SIRP alpha V2, hFc Tag ELISA

Immobilized Human SIRP alpha V2, hFc Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Biotinylated Anti-SIRP alpha Antibody, hFc Tag with the EC50 of 33.9ng/ml determined by ELISA.