

Biotinylated Human SIRP Beta 1 Isoform 3 Protein

Cat. No. SRP-HM4BLB

Description	
Source	Recombinant Biotinylated Human SIRP Beta 1 Isoform 3 Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Glu30-Leu371.
Accession	Q5TFQ8-1
Molecular Weight	The protein has a predicted MW of 39.9 kDa. Due to glycosylation, the protein migrates to 55-68 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC


Formulation and Storage	
Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Signal regulatory protein $\beta 1$ (SIRPB1) is a signal regulatory protein member of the immunoglobulin superfamily and is capable of modulating receptor tyrosine kinase-coupled signaling. Copy number variations at the SIRPB1 locus were previously reported to associate with prostate cancer aggressiveness in patients. SIRPB1 gene amplification and overexpression were detected in prostate cancer specimens. The knockdown of SIRPB1 significantly suppressed cell growth in colony formation assays and cell mobility.

Assay Data

Bis-Tris PAGE



Biotinylated Human SIRP Beta 1 Isoform 3 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Biotinylated Human SIRP Beta 1 Isoform 3 Protein

Cat. No. SRP-HM4BLB



Assay Data



The purity of Biotinylated Human SIRP Beta 1 Isoform 3 is greater than 95% as determined by SEC-HPLC.