Mouse Transferrin R/CD71 Protein

Cat. No. TFR-MM101



Description	
Source	Recombinant Mouse Transferrin R/CD71 Protein is expressed from HEK293 with His tag at the N-Terminus.
	It contains Cys89-Phe763.
Accession	Q62351
Molecular Weight	The protein has a predicted MW of 76.99 kDa. Due to glycosylation, the protein migrates to 78-90 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage


pr	rotectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 μg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Siorage	20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend of aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

The transferrin receptor (CD71) is known as a receptor for IgA1 on mesangial cells. The transcript and protein expression of glomerular CD71 was determined by real-time polymerase chain reaction and immunohistochemistry. Human mesangial cells (HMCs) were treated with sera from IgAN patients and expression levels of CD71 and inflammatory cytokine markers were compared according to disease status.

Assay Data

Bis-Tris PAGE




Mouse Transferrin R on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Cat. No. TFR-MM101



Assay Data



The purity of Mouse Transferrin R is greater than 95% as determined by SEC-HPLC.