

Biotinylated Cynomolgus TIGIT Protein

Cat. No. TIG-CM410B

Description	
Source	Recombinant Biotinylated Cynomolgus TIGIT Protein is expressed from HEK293 with His tag and Avi tag at the C-Terminus. It contains Met89-Ile208.
Accession	XP_015300911.1
Molecular Weight	The protein has a predicted MW of 16.2 kDa. Due to glycosylation, the protein migrates to 20-28 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE


Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

TIGIT, also called Vstm3, Vsig9, and WUCAM, is a transmembrane protein in the CD28 family of the Ig superfamily proteins. IGIT is expressed at low levels on peripheral memory and regulatory CD4 T-cells and NK cells and is up-regulated following activation of these cells.

Assay Data

Bis-Tris PAGE




Biotinylated Cynomolgus TIGIT on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

ELISA Data

Biotinylated Cynomolgus TIGIT, His Tag ELISA

0.05µg Biotinylated Cynomolgus TIGIT, His Tag Per Well



Immobilized Biotinylated Cynomolgus TIGIT, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Anti-TIGIT Antibody, hFc Tag with the EC50 of 59.5ng/ml determined by ELISA.