## **Cynomolgus TIM-4 Protein**


# κλιτυς

# Cat. No. TIM-CM104

| Description             |   |
|-------------------------|---|
| Source                  | Recombinant Cynomolgus TIM-4 Protein is expressed from HEK293 with His tag at the C-Terminus.   |
|                         | It contains Glu25-Gln305.   |
| Accession               | XP_005558436.2  |
| Molecular<br>Weight     | The protein has a predicted MW of 31.55 kDa. Due to glycosylation, the protein migrates to 70-80 kDa based on Bis-Tris PAGE result.   |
| Endotoxin               | Less than 1EU per µg by the LAL method.   |
| Purity                  | > 95% as determined by Bis-Tris PAGE  |
|                         | > 95% as determined by HPLC   |
| Formulation and Storage |   |
| Formulation             | Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.  |
| Reconstitution          | Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.  |
| Storage                 | -20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.   |
| Background              |   |
|                         | Tim4 is a transmembrane protein as the phosphatidylserine (PS) receptor, known as T cell immunoglobulin and mucin domain containing protein-4. It is expressed highly in macrophages, and macrophage Tim-4 inhibits inflammation under various conditions of immune activation. |


#### Assay Data





Cynomolgus TIM-4 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

## SEC-HPLC



The purity of Cynomolgus TIM-4 is greater than 95% as determined by SEC-HPLC.