Mouse TIMP-2 Protein


Cat. No. TIM-MM1P2

ϗͶͼͻ·Ͷϩ

Description	
Source	Recombinant Mouse TIMP-2 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Cys27-Pro220.
Accession	P25785
Molecular Weight	The protein has a predicted MW of 22.8 kDa. Due to glycosylation, the protein migrates to 24-26 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	Tissue inhibitors of metalloproteinases or TIMPs are a family of proteins that regulate the activation and proteolytic activity of the zinc enzymes known as matrix metalloproteinases (MMPs).TIMP-2 is a non N- glycosylated protein with a molecular mass of 22 kDa produced by a wide range of cell types, which inhibits MMPs non-covalently by the formation of binary complexes. TIMP-2 also has erythroidpotentiating and cell growth promoting activities.


Assay Data

Bis-Tris PAGE



Mouse TIMP-2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC



The purity of Mouse TIMP-2 is greater than 95% as determined by SEC-HPLC.