### Human Tenascin Protein


## Cat. No. TNC-HM101

# ϗͶϲϿ·ႮႽ

Description	
Source	Recombinant Human Tenascin Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Gly23-Ser621.
Accession	P24821-1
Molecular Weight	The protein has a predicted MW of 65.27 kDa. Due to glycosylation, the protein migrates to 66-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	Tenascin-C (TNC) is a hexameric, multimodular extracellular matrix protein with several molecular forms that are created through alternative splicing and protein modifications. It is highly conserved amongst vertebrates, and molecular phylogeny indicates that it evolved before fibronectin. Tenascin-C has many extracellular binding partners, including matrix components, soluble factors and pathogens; it also influences cell phenotype directly through interactions with cell surface receptors.


### Assay Data

#### **Bis-Tris PAGE**



10K[





Human Tenascin on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

The purity of Human Tenascin is greater than 95% as determined by SEC-HPLC.