Cynomolgus TNFR2/CD120b/TNFRSF1B Protein





Description	
Source	Recombinant Cynomolgus TNFR2/CD120b/TNFRSF1B Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Leu23-Asp257.
Accession	A0A2K5VEV7
Molecular Weight	The protein has a predicted MW of 26.1 kDa. Due to glycosylation, the protein migrates to 49-60 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC
Formulation and	d Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Tumor Necrosis Factor Receptor II (TNF RII), also known as TNFRSF1B, p75/p80, and CD120b, is a type I transmembrane protein that belongs to the TNF receptor superfamily. It has a molecular weight of approximately 75 kDa.Receptor with high affinity for TNFSF2/TNF-alpha and approximately 5-fold lower affinity for homotrimeric TNFSF1/lymphotoxin-alpha. The TRAF1/TRAF2 complex recruits the apoptotic suppressors BIRC2 and BIRC3 to TNFRSF1B/TNFR2. This receptor mediates most of the metabolic effects of TNF-alpha. Isoform 2 blocks TNF-alpha-induced apoptosis, which suggests that it regulates TNF-alpha function by antagonizing its biological activity.

Assay Data

Bis-Tris PAGE




Cynomolgus TNFR2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS


Assay Data



The purity of Cynomolgus TNFR2 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Cynomolgus TNFR2, His Tag ELISA 0.1µg Cynomolgus TNFR2, His Tag Per Well



Log Anti-TNFR2 Antibody, hFc Tag Conc.(μg/ml)

Immobilized Cynomolgus TNFR2, His Tag at 1 μ g/ml (100 μ l/Well) on the plate. Dose response curve for Anti-TNFR2 Antibody, hFc Tag with the EC50 of 6.0ng/ml determined by ELISA (QC Test).