Ferret TNF alpha Protein

Cat. No. TNF-FE00A



Description	
Source	Recombinant Ferret TNF alpha Protein is expressed from E.coli without tag.
	It contains Val77-Leu233.
Accession	A3FBF1
Molecular Weight	The protein has a predicted MW of 17.21 kDa same as Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Tumor necrosis factor alpha (TNF-alpha), also known as cachectin and TNFSF2, is the prototypic ligand of the TNF superfamily. It is a pleiotropic molecule that plays a central role in inflammation, immune system development, apoptosis, and lipid metabolism. Cytokine that binds to TNFRSF1A/TNFR1 and TNFRSF1B/TNFBR. It is mainly secreted by macrophages and can induce cell death of certain tumor cell lines. It is potent pyrogen causing fever by direct action or by stimulation of interleukin-1 secretion and is implicated in the induction of cachexia, Under certain conditions it can stimulate cell proliferation and induce cell differentiation.

Assay Data

Tris-Bis PAGE




Ferret TNF alpha on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

KAGTUS

Assay Data




The purity of Ferret TNF alpha is greater than 95% as determined by SEC-HPLC.

ELISA Data

Ferret TNF-alpha, No Tag ELISA

0.05μg Ferret TNF-alpha, No Tag Per Well



Immobilized Ferret TNF-alpha, No Tag at $0.5\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Human TNFR2, hFc Tag with the EC50 of 14.5ng/ml determined by ELISA.