

Human TNFR1/CD120a/TNFRSF1A Protein

Cat. No. TNF-HM2R1

Description

Source	Recombinant Human TNFR1/CD120a/TNFRSF1A Protein is expressed from Expi293 with hFc tag at the C-terminal. It contains Leu30-Thr211.
Accession	P19438-1
Molecular Weight	The protein has a predicted MW of 48 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 5% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge tubes before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -20 to -80°C for 3-6 months in unopened state after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please avoid freeze-thaw cycles.

Background

Tumour necrosis factor alpha (TNF-α) is a pleiotropic cytokine with both injurious and protective functions, which are thought to diverge at the level of its two cell surface receptors, TNFR1 and TNFR2. In the setting of acute injury, selective inhibition of TNFR1 is predicted to attenuate the cell death and inflammation associated with TNF-α, while sparing or potentiating the protective effects of TNFR2 signalling.

Assay Data


Tris-Bis PAGE



Human TNFR1 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC

Assay Data




The purity of Human TNFR1 is greater than 95% as determined by SEC-HPLC.

ELISA Data

Human TNFR1, hFc Tag ELISA

0.05µg Human TNF alpha, His Tag Per Well



Immobilized Human TNF alpha, His Tag at 0.5µg/ml (100µl/Well) on the plate. Dose response curve for Human TNFR1, hFc Tag with the EC50 of 8.7ng/ml determined by ELISA.