

Biotinylated Cynomolgus TRAIL R2/DR5/TNFRSF10B Protein (Primary Amine Labeling)



Cat. No. TRL-CM1R2B

Description

Source	Recombinant Biotinylated Cynomolgus TRAIL R2/DR5/TNFRSF10B Protein (Primary Amine Labeling) is expressed from HEK293 with His tag at the C-terminus. It contains Ile58-Ser212.
Accession	A0A2K5TXK0
Molecular Weight	The protein has a predicted MW of 18.28 kDa. Due to glycosylation, the protein migrates to 30-38 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1 EU per µg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage


Formulation	Lyophilized from 0.22 µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

DR5, also called TRAIL R2, TRICK 2, TNFRSF10B, and MK is a type 1 TNF R superfamily, membrane protein which is a receptor for TRAIL (APO2 ligand). DR5 is a receptor for the cytotoxic ligand TNFSF10/TRAIL. The resulting death-inducing signaling complex (DISC) performs caspase-8 proteolytic activation which initiates the subsequent cascade of caspases (aspartate-specific cysteine proteases) mediating apoptosis.


Assay Data

Bis-Tris PAGE



Biotinylated Cynomolgus TRAIL R2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Biotinylated Cynomolgus TRAIL R2 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Biotinylated Cynomolgus TRAIL R2, His Tag ELISA

0.05µg Biotinylated Cynomolgus TRAIL R2, His Tag Per Well




Immobilized Biotinylated Cynomolgus TRAIL R2, His Tag at 0.5µg/ml (100µl/Well) on streptavidin (5µg/ml) precoated plate. Dose response curve for Anti-TRAIL R2 Antibody, hFc Tag with the EC50 of 3.5ng/ml determined by ELISA (QC Test).

ELISA Data

Biotinylated Cynomolgus TRAIL R2, His Tag ELISA

0.5µg Human TRAIL, No Tag Per Well



Immobilized Human TRAIL, No Tag at 5µg/ml(100µl/well) on the plate. Dose response curve for Biotinylated Cynomolgus TRAIL R2, His Tag with the EC50 of 73.7ng/ml determined by ELISA.