

Human TROP-2/TACSTD2 Protein

Cat. No. TRP-HM221

Description

Source	Recombinant Human TROP-2/TACSTD2 Protein is expressed from HEK293 with hFc tag at the C-Terminus. It contains His27-Thr274.
Accession	P09758
Molecular Weight	The protein has a predicted MW of 54.6 kDa. Due to glycosylation, the protein migrates to 60-70 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22 μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 $\mu\text{g}/\text{ml}$ is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt. -80°C for 3 months after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Trop-2, also known as epithelial glycoprotein-1 antigen (EGP-1), is a protein that in humans is encoded by the TACSTD2 gene. Mutations of this gene result in gelatinous drop-like corneal dystrophy, an autosomal recessive disorder characterized by severe corneal amyloidosis leading to blindness.


Assay Data

Bis-Tris PAGE



Human TROP-2 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.

SEC-HPLC




The purity of Human TROP-2 is greater than 95% as determined by SEC-HPLC.

Assay Data

ELISA Data

Human TROP-2, hFc Tag ELISA

0.1µg Human TROP-2, hFc Tag Per Well



Immobilized Human TROP-2, hFc Tag at 1µg/ml (100µl/well) on the plate. Dose response curve for Biotinylated Anti-TROP-2 Antibody, hFc Tag with the EC50 of 6.7ng/ml determined by ELISA.