Biotinylated Cynomolgus TSLP Protein (Primary Amine Labeling)






Description	
Source	Recombinant Biotinylated Cynomolgus TSLP Protein (Primary Amine Labeling) is expressed from HEK293 with His tag at the C-Terminus.
	It contains Tyr29-Gln159.
Accession	A0A7N9CAT7
Molecular Weight	The protein has a predicted MW of 16.3 kDa. Due to furin cleavage site, the protein migrates to 10-13 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 90% as determined by Bis-Tris PAGE
	> 90% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22 μ m filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt80°C for 3-6 months after reconstitution. 2-8°C for 2-7 days after reconstitution. Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	

Thymic Stromal Lymphopoietin (TSLP) was originally identified as an activity from the conditioned medium of a mouse thymic stromal cell line that promoted the development of B cells. Cytokine that induces the release of T-cell-attracting chemokines from monocytes and, in particular, enhances the maturation of CD11c dendritic cells. Can induce allergic inflammation by directly activating mast cells.

Assay Data

Bis-Tris PAGE




Biotinylated Cynomolgus TSLP on Bis-Tris PAGE under reduced conditions. The purity is greater than 90%.

SEC-HPLC

ΚΛ**ω**τυς

Assay Data




The purity of Biotinylated Cynomolgus TSLP is greater than 90% as determined by SEC-HPLC.

ELISA Data

Biotinylated Cynomolgus TSLP, His Tag ELISA

0.05μg Biotinylated Cynomolgus TSLP, His Tag Per Well




Immobilized Biotinylated Cynomolgus TSLP, His Tag at 0.5µg/ml (100µl/well) on the streptavidin precoated plate (5µg/ml). Dose response curve for Anti-TSLP Antibody, hFc Tag with the EC50 of 11.2ng/ml determined by ELISA (QC Test).

ELISA Data

Biotinylated Cynomolgus TSLP, His Tag ELISA

0.2μg Cynomolgus TSLPR, His Tag Per Well



Log Biotinylated Cynomolgus TSLP, His Tag Conc.(μg/ml)

Immobilized Cynomolgus TSLPR, His Tag at $2\mu g/ml$ (100 $\mu l/well$) on the plate. Dose response curve for Biotinylated Cynomolgus TSLP, His Tag with the EC50 of 0.12 $\mu g/ml$ determined by ELISA.