Human TSPAN8 Protein

Cat. No. TSP-HM2N8

ϗͶϲϿ·ႮႽ

Description	
Source	Recombinant Human TSPAN8 Protein is expressed from HEK293 with hFc tag at the N-Terminus.
	It contains Lys110-Asn205.
Accession	P19075
Molecular Weight	The protein has a predicted MW of 37.30 kDa. Due to glycosylation, the protein migrates to 43-48 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per µg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC
Formulation and Storage	
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.
Background	
	Tetraspanin 8 (TSPAN8) is a member of the tetraspanin superfamily that forms TSPAN8-mediated protein complexes by interacting with themselves and other various cellular signaling molecules. In physiological conditions, TSPAN8 plays a vital role in the regulation of biological functions, including leukocyte trafficking, angiogenesis and wound repair.

Assay Data

Tris-Bis PAGE



Human TSPAN8 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

Human TSPAN8 Protein

Cat. No. TSP-HM2N8

Assay Data





The purity of Human TSPAN8 is greater than 95% as determined by SEC-HPLC.