

Human VCAM-1/CD106 Protein

Cat. No. VAM-HM106

Description

| | |
|-------------------------|---|
| Source | Recombinant Human VCAM-1/CD106 Protein is expressed from HEK293 with His tag at the C-Terminus. It contains Phe25-Pro697. |
| Accession | NP_001069 |
| Molecular Weight | The protein has a predicted MW of 75.2 kDa. Due to glycosylation, the protein migrates to 75-110 kDa based on Bis-Tris PAGE result. |
| Endotoxin | Less than 1EU per μg by the LAL method. |
| Purity | > 95% as determined by Bis-Tris PAGE > 95% as determined by HPLC |

Formulation and Storage

| | |
|--------------------|---|
| Formulation | Supplied as 0.22 μm filtered solution in PBS (pH 7.4). |
| Storage | Valid for 12 months from date of receipt when stored at -80°C . Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles. |

Background

Tumor necrosis factor alpha (TNF α) is a pro-inflammatory cytokine that triggers the expression of inflammatory molecules, including other cytokines and cell adhesion molecules. TNF α induces the expression of intercellular cell adhesion molecule-1 and vascular cell adhesion molecule-1 (VCAM-1). VCAM-1 was originally identified as a cell adhesion molecule that helps regulate inflammation-associated vascular adhesion and the transendothelial migration of leukocytes, such as macrophages and T cells.

Assay Data

Bis-Tris PAGE



Human VCAM-1 on Bis-Tris PAGE under reduced condition. The purity is greater than 95%.


SEC-HPLC

Human VCAM-1/CD106 Protein

Cat. No. VAM-HM106



Assay Data



The purity of Human VCAM-1 is greater than 95% as determined by SEC-HPLC.