Human VEGF165 Protein

Cat. No. VEG-HM065

Description

Source	Recombinant Human VEGF165 Protein is expressed from HEK293 without tag.
	It contains Ala27-Arg191.
Accession	P15692-4
Molecular Weight	The protein has a predicted MW of 19.2 kDa. Due to glycosylation, the protein migrates to 20-30 kDa based on Bis-Tris PAGE result.
Endotoxin	Less than 0.05 EU per μg by the LAL method.
Purity	> 95% as determined by Bis-Tris PAGE
	> 95% as determined by HPLC

Formulation and Storage

Formulation	Lyophilized from 0.22μm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.
Reconstitution	Dissolve the lyophilized protein in distilled water. Please refer to the Certificate of Analysis for detailed instructions.
Storage	-20 to -80°C for 12 months as supplied from date of receipt.-80°C for 3 months after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.

Background

Human papillomavirus (HPV) infection is an established risk factor for cervical carcinogenesis. VEGF165 was significantly higher, whereas VEGFC and VEGFD were significantly lower in malignant cervical carcinoma tissues as compared to normal cervix tissues. Expression levels of VEGF121 and VEGFC were significantly associated with type of tumor growth while VEGF165 was significantly associated with lymph node metastasis.

Assay Data

Bis-Tris PAGE



Human VEGF165 on Bis-Tris PAGE under reduced condition and Non reduced (N) condition. The purity is greater than 95%.

SEC-HPLC



The purity of Human VEGF165 is greater than 95% as determined by SEC-HPLC.

Human VEGF165 Protein

Cat. No. VEG-HM065

Assay Data

Cell Based Assay



The bioactivity was measured using a luciferase reporter cell line which stably expresses the KDR receptor. The ED50 for this effect is < 10 ng/mL.

ELISA Data



Immobilized Human VEGF165 at 1μg/ml (100μl/well) on the plate. Dose response curve for Biotinylated Human VEGFR1, His Tag with the EC50 of 46.4ng/ml determined by ELISA.

ELISA Data



Immobilized Human VEGF165, No Tag at 5μg/ml (100μl/well) on the plate. Dose response curve for Human Neuropilin-1, His Tag (Cat. NRP-HM101) with the EC50 of 1.26μg/ml determined by ELISA.

ELISA Data

Human VEGF165 Protein

Cat. No. VEG-HM065

Assay Data



Immobilized Human VEGF165, No Tag at 1μg/ml (100μl/well) on the plate. Dose response curve for Anti-VEGF165 Antibody, hFc Tag with the EC50 of 2.7ng/ml determined by ELISA.

Blocking Data



Serial dilutions of Anti-VEGF165 Antibody were added into Human VEGF165, No Tag: Biotinylated Human VEGF R2, His Avi Tag binding reactioins. The half maximal inhibitiory concentration (IC50) is 0.30μg/ml.

For Research Use Only