Cynomolgus VSIG4 Protein

Cat. No. VSG-CM104



Description	
Source	Recombinant Cynomolgus VSIG4 Protein is expressed from HEK293 with His tag at the C-Terminus.
	It contains Arg25-Ala288.
Accession	A0A7N9D9P6
Molecular Weight	The protein has a predicted MW of 30.16 kDa. Due to glycosylation, the protein migrates to 35-50 kDa based on Tris-Bis PAGE result.
Endotoxin	Less than 1EU per μg by the LAL method.
Purity	> 95% as determined by Tris-Bis PAGE
	> 95% as determined by HPLC

Formulation and Storage


Formulation and Storage		
Formulation	Lyophilized from 0.22µm filtered solution in PBS (pH 7.4). Normally 8% trehalose is added as protectant before lyophilization.	
Reconstitution	Centrifuge the tube before opening. Reconstituting to a concentration more than 100 µg/ml is recommended. Dissolve the lyophilized protein in distilled water.	
Storage	-20 to -80°C for 12 months as supplied from date of receipt20 to -80°C for 3-6 months in unopened state after reconstitution.2-8°C for 2-7 days after reconstitution.Recommend to aliquot the protein into smaller quantities for optimal storage. Please minimize freeze-thaw cycles.	

Background

VSIG4, a B7 family-related protein, is a negative regulator of T cell activation. T cell activation by APCs is positively and negatively regulated by members of the B7 family. Unlike that of B7 family members, surface expression of VSIG4 was restricted to resting tissue macrophages and absent upon activation by LPS or in autoimmune inflammatory foci. The specific expression of VSIG4 on resting macrophages in tissue suggests that this inhibitory ligand may be important for the maintenance of T cell unresponsiveness in healthy tissues.

Assay Data

Tris-Bis PAGE




SEC-HPLC

Cynomolgus VSIG4 on Tris-Bis PAGE under reduced condition. The purity is greater than 95%.

Cat. No. VSG-CM104



Assay Data



The purity of Cynomolgus VSIG4 is greater than 95% as determined by SEC-HPLC.